

# SOME INTEGRALS INVOLVING GENERALIZED $\mathbf{k}$ -STRUVE FUNCTIONS.

KOTTAKKARAN SOOPPY NISAR AND SAIFUL RAHMAN MONDAL

ABSTRACT. The close form of some integrals involving recently developed generalized  $\mathbf{k}$ -Struve functions is obtained. The outcome of these integrations is expressed in terms of generalized Wright functions. Several special cases are deduced which lead to some known results.

## 1. INTRODUCTION AND PRELIMINARIES

The Struve function of order  $p$  given by

$$\mathbf{H}_p(x) := \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(k + 3/2) \Gamma(k + p + \frac{3}{2})} \left(\frac{x}{2}\right)^{2k+p+1}, \quad (1.1)$$

is a particular solution of the non-homogeneous Bessel differential equation

$$x^2 y''(x) + xy'(x) + (x^2 - p^2)y(x) = \frac{4 \left(\frac{x}{2}\right)^{p+1}}{\sqrt{\pi} \Gamma(p + 1/2)}. \quad (1.2)$$

Recently, Nisar et. al, [16] introduced and studied various properties of  $\mathbf{k}$ -Struve function  $\mathbf{S}_{\nu, c}^{\mathbf{k}}$  defined by

$$\mathbf{S}_{\nu, c}^{\mathbf{k}}(x) := \sum_{r=0}^{\infty} \frac{(-c)^r}{\Gamma_{\mathbf{k}}(r\mathbf{k} + \nu + \frac{3\mathbf{k}}{2}) \Gamma(r + \frac{3}{2})} \left(\frac{x}{2}\right)^{2r + \frac{\nu}{\mathbf{k}} + 1}. \quad (1.3)$$

The generalized Wright hypergeometric function  ${}_p\psi_q(z)$  is given by the series

$${}_p\psi_q(z) = {}_p\psi_q \left[ \begin{matrix} (a_i, \alpha_i)_{1,p} \\ (b_j, \beta_j)_{1,q} \end{matrix} \middle| z \right] = \sum_{k=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + \alpha_i k)}{\prod_{j=1}^q \Gamma(b_j + \beta_j k)} \frac{z^k}{k!}, \quad (1.4)$$

where  $a_i, b_j \in \mathbb{C}$ , and real  $\alpha_i, \beta_j \in \mathbb{R}$  ( $i = 1, 2, \dots, p; j = 1, 2, \dots, q$ ). Asymptotic behavior of this function for large values of argument of  $z \in \mathbb{C}$  were studied in [7] and under the condition

$$\sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > -1 \quad (1.5)$$

was found in the work of [18, 19]. Properties of this generalized Wright function were investigated in [9], (see also [8, 10]). In particular, it was proved [9] that  ${}_p\psi_q(z)$ ,  $z \in \mathbb{C}$  is an entire function under the condition (1.5).

---

2010 *Mathematics Subject Classification.* 33B20, 33C20, 33B15, 33C05.

*Key words and phrases.* gamma function,  $k$ -gamma function, Lavoie-Trottier integral formula, Wright function,  $\mathbf{k}$ -Struve function.

The  $k$ -gamma function defined in [4] as:

$$\Gamma_k(z) = \int_0^\infty t^{z-1} e^{-\frac{t^k}{k}} dt, z \in \mathbb{C}. \quad (1.6)$$

By inspection the following relation holds:

$$\Gamma_k(z+k) = z\Gamma_k(z) \quad (1.7)$$

and

$$\Gamma_k(z) = k^{\frac{z}{k}-1} \Gamma\left(\frac{z}{k}\right). \quad (1.8)$$

If  $k \rightarrow 1$  and  $c = 1$ , then the generalized  $k$ -Bessel function defined in (2.1) reduces to the well known classical Bessel function  $J_\nu$  defined in [6].

In this paper, we define a class of integral formulas which involves the generalized  $\mathbf{k}$ -Struve function as defined in (1.3). Also, we investigate some special cases as the corollaries. For this purpose, we recall the following result of Lavoie and Trottier [12].

$$\begin{aligned} & \int_0^1 x^{\alpha-1} (1-x)^{2\beta-1} \left(1 - \frac{x}{3}\right)^{2\alpha-1} \left(1 - \frac{x}{4}\right)^{\beta-1} dx \\ &= \left(\frac{2}{3}\right)^{2\alpha} \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \Re(\alpha) > 0, \Re(\beta) > 0. \end{aligned} \quad (1.9)$$

For more details about integral representations of various special function, the reader can refer these papers (see [2], [3], [13], [14], [15]).

## 2. MAIN RESULT

In this section, we establish two generalized integral formulas containing (1.3), which represented in terms of generalized Wright function defined in (1.4) by inserting with the suitable argument defined in (1.9).

**Theorem 2.1.** For  $\mathbf{k} \in \mathbb{R}, \alpha, \mu, \beta, \nu, c \in \mathbb{C}$  with  $\nu > \frac{3}{2}\mathbf{k}, \Re(\alpha + \mu) > 0, \Re(\alpha + \frac{\nu}{\mathbf{k}} + 1) > 0$  and  $x > 0$ , then the following integral formula hold true:

$$\begin{aligned} & \int_0^1 x^{\alpha+\mu-1} (1-x)^{2\alpha-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{4}\right)^{\alpha-1} \mathbf{S}_{\nu,c}^{\mathbf{k}} \left( \frac{y \left(1 - \frac{x}{4}\right) (1-x)^2}{2} \right) dx \\ &= \frac{\left(\frac{y}{2}\right)^{\frac{\nu}{\mathbf{k}}+1} \Gamma(\alpha + \mu) \left(\frac{2}{3}\right)^{2(\alpha+\mu)}}{k^{\frac{\nu}{\mathbf{k}}}} \\ & \times {}_2\Psi_3 \left[ \begin{matrix} (\alpha + \frac{\nu}{\mathbf{k}} + 1, 2), (1, 1); \\ (\frac{\nu}{\mathbf{k}} + \frac{3}{2}, 1), (\frac{3}{2}, 1), (2\alpha + \frac{\nu}{\mathbf{k}} + \mu, 2) \end{matrix} \middle| -\frac{cy^2}{4\mathbf{k}} \right]. \end{aligned} \quad (2.1)$$

*Proof.* Let  $I$  be the left hand side of (2.1) and applying (1.3) to the integrand of (2.1), we have

$$\begin{aligned} & \int_0^1 x^{\alpha+\mu-1} (1-x)^{2\alpha-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{4}\right)^{\alpha-1} \mathbf{S}_{\nu,c}^{\mathbf{k}} \left( \frac{y \left(1 - \frac{x}{4}\right) (1-x)^2}{2} \right) dx \\ &= \int_0^1 x^{\alpha+\mu-1} (1-x)^{2\alpha-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{4}\right)^{\alpha-1} \\ & \times \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{y}{2}\right)^{2r+\frac{\nu}{\mathbf{k}+1}} \left(1 - \frac{x}{4}\right)^{2r+\frac{\nu}{\mathbf{k}}} (1-x)^{2(2r+\frac{\nu}{\mathbf{k}+1})}}{\Gamma_{\mathbf{k}}(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} dx \end{aligned}$$

Interchanging the order of integration and summation, which is verified by the uniform convergence of the series under the given assumption of theorem 2.1, we have

$$\begin{aligned} I &= \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{y}{2}\right)^{2r+\frac{\nu}{\mathbf{k}}+1}}{\Gamma_{\mathbf{k}}(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} \\ & \times \int_0^1 x^{\alpha+\mu-1} (1-x)^{2(2r+\alpha+\frac{\nu}{\mathbf{k}}+1)-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\frac{\nu}{\mathbf{k}}+1+2r)-1} dx, \end{aligned}$$

From the assumption given in theorem 2.1, since  $\Re(\frac{\nu}{\mathbf{k}}) > 0$ ,  $\Re(\alpha + \frac{\nu}{\mathbf{k}} + 1 + 2r) > \Re(\alpha + \frac{\nu}{\mathbf{k}} + 1) > 0$ ,  $\Re(\alpha + \mu) > 0$ ,  $\mathbf{k} > 0$  and using (1.9), we get

$$I = \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{y}{2}\right)^{2r+\frac{\nu}{\mathbf{k}}+1}}{\Gamma_{\mathbf{k}}(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} \left(\frac{2}{3}\right)^{2(\alpha+\mu)} \frac{\Gamma(\alpha + \mu) \Gamma(2r + \alpha + \frac{\nu}{\mathbf{k}} + 1)}{\Gamma(2r + 2\alpha + \mu + \frac{\nu}{\mathbf{k}} + 1)}$$

Using (1.8), we get

$$I = \left(\frac{2}{3}\right)^{2(\alpha+\mu)} \frac{\Gamma(\alpha + \mu) \left(\frac{y}{2}\right)^{\frac{\nu}{\mathbf{k}}}}{\mathbf{k}^{\frac{\nu}{\mathbf{k}}+\frac{1}{2}}} \frac{\Gamma(2r + \alpha + \frac{\nu}{\mathbf{k}} + 1)}{\Gamma(2r + 2\alpha + \mu + \frac{\nu}{\mathbf{k}} + 1) \Gamma_{\mathbf{k}}(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})}$$

In view of definition of Fox-Wright function (1.4), we get the required result.  $\square$

**Theorem 2.2.** For  $\mathbf{k} \in \mathbb{R}$ ,  $\alpha, \mu, \beta, \nu, c \in \mathbb{C}$  with  $\nu > \frac{3}{2}\mathbf{k}$ ,  $\Re(\alpha + \mu) > 0$ ,  $\Re(\alpha + \frac{\nu}{\mathbf{k}} + 1) > 0$  and  $x > 0$ , then the following integral formula hold true:

$$\begin{aligned} & \int_0^1 x^{\alpha-1} (1-x)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{3}\right)^{2\alpha-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\mu)-1} \mathbf{S}_{\nu,c}^{\mathbf{k}} \left( \frac{yx \left(1 - \frac{x}{3}\right)^2}{2} \right) dx \\ &= \frac{\left(\frac{y}{2}\right)^{\frac{\nu}{\mathbf{k}}+1} \Gamma(\alpha + \mu) \left(\frac{2}{3}\right)^{2\alpha}}{\mathbf{k}^{\frac{\nu}{\mathbf{k}}} 3^{\frac{2\nu}{\mathbf{k}}+2}} \\ & \times {}_2\Psi_3 \left[ \begin{matrix} (\alpha + \frac{\nu}{\mathbf{k}} + 1, 2), (1, 1); \\ (\frac{\nu}{\mathbf{k}} + \frac{3}{2}, 1), (\frac{3}{2}, 1), (2\alpha + \frac{\nu}{\mathbf{k}} + \mu, 2) \end{matrix} \middle| -\frac{cy^2}{4\mathbf{k}} \right]. \end{aligned} \quad (2.2)$$

*Proof.* Let  $\ell$  be the left hand side of (2.2) and applying (1.3) to the integrand of (2.1), we have

$$\begin{aligned} & \int_0^1 x^{\alpha-1} (1-x)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{3}\right)^{2\alpha-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\mu)-1} \mathbf{S}_{\nu,c}^{\mathbf{k}} \left( \frac{yx \left(1 - \frac{x}{3}\right)^2}{2} \right) dx \\ &= \int_0^1 x^{\alpha-1} (1-x)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{3}\right)^{2\alpha-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\mu)-1} \\ & \times \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{xy}{2}\right)^{2r+\frac{\nu}{\mathbf{k}+1}} \left(1 - \frac{x}{3}\right)^{2(2r+\frac{\nu}{\mathbf{k}+1})}}{\Gamma_k(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} dx \end{aligned}$$

Interchanging the order of integration and summation, which is verified by the uniform convergence of the series under the given assumption of theorem 2.2, we have

$$\begin{aligned} \ell &= \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{y}{2}\right)^{2r+\frac{\nu}{\mathbf{k}+1}}}{\Gamma_k(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} \\ & \times \int_0^1 x^{(\alpha+\frac{\nu}{\mathbf{k}+1}+2r)-1} (1-x)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mathbf{k}+1+2r)-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\mu)-1} dx, \end{aligned}$$

Since  $\Re(\frac{\nu}{\mathbf{k}}) > 0$ ,  $\Re(\alpha + \frac{\nu}{\mathbf{k}} + 1 + 2r) > \Re(\alpha + \frac{\nu}{\mathbf{k}} + 1) > 0$ ,  $\Re(\alpha + \mu) > 0$ ,  $\mathbf{k} > 0$  and using (1.9), we get

$$\ell = \sum_{r=0}^{\infty} \frac{(-c)^r \left(\frac{y}{2}\right)^{2r+\frac{\nu}{\mathbf{k}+1}}}{\Gamma_k(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})} \left(\frac{2}{3}\right)^{2(\alpha+\frac{\nu}{\mathbf{k}+1}+2r)} \frac{\Gamma(\alpha + \mu) \Gamma(2r + \alpha + \frac{\nu}{\mathbf{k}} + 1)}{\Gamma(2r + 2\alpha + \mu + \frac{\nu}{\mathbf{k}} + 1)}$$

Using (1.8), we obtain

$$\ell = \left(\frac{2}{3}\right)^{2(\alpha+\frac{\nu}{\mathbf{k}+1})} \frac{\Gamma(\alpha + \mu) \left(\frac{y}{2}\right)^{\frac{\nu}{\mathbf{k}+1}}}{\mathbf{k}^{\frac{\nu}{\mathbf{k}}+\frac{1}{2}}} \frac{\Gamma(2r + \alpha + \frac{\nu}{\mathbf{k}} + 1)}{\Gamma(2r + 2\alpha + \mu + \frac{\nu}{\mathbf{k}} + 1) \Gamma_k(r\mathbf{k} + \nu + \frac{3}{2}\mathbf{k}) \Gamma(r + \frac{3}{2})}$$

In view of definition of Fox-Wright function (1.4), we get the desired result.  $\square$

### 3. SPECIAL CASES

In this section, we obtain the integral representation of Struve function and modified Struve function.

**Case 1.** If we set  $c = \mathbf{k} = 1$  in (1.3), then we get Struve function of order  $\nu$  as

$$\mathbf{H}_{\nu}(x) := \sum_{r=0}^{\infty} \frac{(-1)^r}{\Gamma(r + \nu + \frac{3}{2}) \Gamma(r + \frac{3}{2})} \left(\frac{x}{2}\right)^{2r+\nu+1}. \quad (3.1)$$

**Case 2.** If we set  $c = -1$  and  $\mathbf{k} = 1$  in (1.3), then we get modified Struve function,

$$\mathbf{L}_{\nu}(x) := \sum_{r=0}^{\infty} \frac{1}{\Gamma(r + \nu + \frac{3}{2}) \Gamma(r + \frac{3}{2})} \left(\frac{x}{2}\right)^{2r+\nu+1}. \quad (3.2)$$

**Corollary 3.1.** *Suppose that the conditions of Theorem 2.1 are satisfied. Then the following integral formula holds:*

$$\begin{aligned} & \int_0^1 x^{\alpha+\mu-1} (1-x)^{2\alpha-1} \left(1 - \frac{x}{3}\right)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{4}\right)^{\alpha-1} \mathbf{H}_\nu \left( \frac{y \left(1 - \frac{x}{4}\right) (1-x)^2}{2} \right) dx \\ &= \left(\frac{y}{2}\right)^{\nu+1} \Gamma(\alpha + \mu) \left(\frac{2}{3}\right)^{2(\alpha+\mu)} \\ & \times {}_2\Psi_3 \left[ \begin{matrix} (\alpha + \nu + 1, 2), (1, 1); \\ (\nu + \frac{3}{2}, 1), (\frac{3}{2}, 1), (2\alpha + \nu + \mu, 2) \end{matrix} \middle| -\frac{y^2}{4} \right]. \end{aligned} \quad (3.3)$$

**Corollary 3.2.** *Assume that the conditions of theorem 2.2 satisfied, then the following integral formula hold true:*

$$\begin{aligned} & \int_0^1 x^{\alpha-1} (1-x)^{2(\alpha+\mu)-1} \left(1 - \frac{x}{3}\right)^{2\alpha-1} \left(1 - \frac{x}{4}\right)^{(\alpha+\mu)-1} \mathbf{L}_\nu \left( \frac{yx \left(1 - \frac{x}{3}\right)^2}{2} \right) dx \\ &= \frac{\left(\frac{y}{2}\right)^{\nu+1} \Gamma(\alpha + \mu) \left(\frac{2}{3}\right)^{2\alpha}}{3^{2\nu+2}} \\ & \times {}_2\Psi_3 \left[ \begin{matrix} (\alpha + \nu + 1, 2), (1, 1); \\ (\nu + \frac{3}{2}, 1), (\frac{3}{2}, 1), (2\alpha + \nu + \mu, 2) \end{matrix} \middle| \frac{y^2}{4} \right]. \end{aligned} \quad (3.4)$$

#### REFERENCES

- [1] P. Agarwal , S. Jain, S. Agarwal, M. Nagpal, *On a new class of integrals involving Bessel functions of the first kind*, Communications in Numerical Analysis **2014** (2014) 1–7.
- [2] J. Choi and P. Agarwal, *Certain unified integrals associated with Bessel functions*, Boundary Value Problems **2013** (2013) 95.
- [3] J. Choi, P. Agarwal, S. Mathur and S.D. Purohit, *Certain new integral formulas involving the generalized Bessel functions*, Bull. Korean Math. Soc. **51** (2014) 995–1003.
- [4] R. Dáaz and E. Pariguan, *On hypergeometric functions and Pochhammer k-symbol*, Divulg. Mat. **15 2** (2007) 179–192.
- [5] A. Erdélyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi, *Higher Transcendental Functions*, Vol.1, McGraw-Hill, New York, Toronto, London, 1953.
- [6] A. Erdélyi, W. Magnus, F. *Higher Transcendental Functions*, Vol.2, McGraw-Hill, New York, Toronto, London, 1953.
- [7] C. Fox, *The asymptotic expansion of generalized hypergeometric functions*, Proc. London. Math. Soc., 27(4) (1928), 389–400.
- [8] A. A. Kilbas and N. Sebastian, *Fractional integration of the product of Bessel function of the first kind*, Fract. Calc. Appl. Anal. **13** (2010), no. 2, 159–175.

- [9] A. A. Kilbas, M. Saigo and J. J. Trujillo, On the generalized Wright function, *Fract. Calc. Appl. Anal.* 5(4) (2002), 437–460.
- [10] A. A. Kilbas and N. Sebastian, Generalized fractional integration of Bessel function of the first kind, *Integral Transforms Spec. Funct.* 19 (11-12) (2008), 869–883.
- [11] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies 204, Elsevier, Amsterdam, 2006.
- [12] J. L. Lavoie and G. Trottier, *On the sum of certain Appells series*, *Ganita* **20** 1 (1969) 31-32.
- [13] N. Menaria, S. D. Purohit and R. K. Parmar, *On a new class of integrals involving generalized Mittag-Leffler function*, *Surveys in Mathematics and its Applications* **11** (2016) 1-9.
- [14] K. S. Nisar and S. R. Mondal, *Certain unified integral formulas involving the generalized modified  $k$ -Bessel function of first kind*, arXiv: 1601.06487 [math.CA].
- [15] K. S. Nisar, P. Agarwal and S. Jain, *Some unified integrals associated with Bessel-Struve kernel function*, arXiv:1602.01496v1 [math.CA].
- [16] K S Nisar, S R Mondal and J Choi, *Certain inequalities involving the  $k$ -Struve function*, submitted, 2016
- [17] E. D. Rainville, *Special functions*, Macmillan, New York, 1960.
- [18] E. M. Wright, The asymptotic expansion of integral functions defined by Taylor series, *Philos. Trans. Roy. Soc. London, Ser. A.* 238 (1940), 423–451.
- [19] E. M. Wright, The asymptotic expansion of the generalized hypergeometric function, *Proc. London Math. Soc.* (2) 46(1940), 389–408.

DEPARTMENT OF MATHEMATICS, COLLEGE OF ARTS AND SCIENCE-WADI AL DAWSER, 11991,,  
 PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY, SAUDI ARABIA  
*E-mail address:* ksnisar1@gmail.com, n.sooppy@psau.edu.sa

DEPARTMENT OF MATHEMATICS, KING FAISAL UNIVERSITY, AL AHSA 31982, SAUDI ARABIA  
*E-mail address:* smondal@kfu.edu.sa